

LAROSE, MARIE-PIER

1 Impact of a social skills program on children's stress: A cluster randomized trial

2 Marie-Pier Larose ^a; marie-pier.larose.1@umontreal.ca;

3 Isabelle Ouellet-Morin ^{a-b}; isabelle.ouellet-morin@umontreal.ca;

4 Frank Vitaro ^a; frank.vitaro@umontreal.ca;

5 Marie Claude Geoffroy ^c; marie-claude.geoffroy@mcgill.ca;

6 Marilyn Ahun ^a; marilyn.ahun@umontreal.ca;

7 Richard E. Tremblay ^a; richard.ernest.tremblay@umontreal.ca;

8 Sylvana M. Côté ^{a-d}; sylvana.cote.1@umontreal.ca.

9

10 a. University of Montreal, Canada.

11 b. Research Centre, Montreal Mental Health University Institute (*Institut universitaire de*
12 *santé mentale de Montréal*), Canada.

13 c. McGill University, Montreal, Canada.

14 d. University of Bordeaux, INSERM U1219, Bordeaux, France

15 **Corresponding author:**

16 Sylvana M. Côté

17 University of Montreal

18 3050 Edouard-Montpetit

19 Montreal, QC

20 H3T 1J7

21 Canada

22 sylvana.cote.1@umontreal.ca

23

24 **Author contributions**

25 SMC, IOM, FV, MPL, MCG, and RET conceived and designed the study. SMC, MPL, IOM, and
26 MA drafted the manuscript. RET, FV, MCG and IOM reviewed the manuscript. All authors read
27 and approved the final manuscript after revising it critically for important intellectual content. All
28 authors agreed to be accountable for all aspects of the work.

29

1 **Abstract**

2 **Background:** Most preschool children in Western industrialized countries attend child care
3 during the day while parents work. Studies suggest that child care may be stressful to young
4 children, perhaps because they still lack the social skills to interact daily in a group setting away
5 from parents. This gap in social abilities may be greater for children in lower-income families,
6 who may face more adversity at home, with fewer resources and more social isolation.

7 **Methods:** We conducted a cluster-randomized controlled trial in 2013-2014 to test whether a
8 social skills intervention led by early childhood educators within the child care center could
9 reduce diurnal cortisol levels to more typical patterns expected of children this age. We
10 randomized 19 public child care centers ($n = 361$ children) in low-income neighborhoods of
11 Montreal, Canada, to either: 1) the Minipally program – intervention group ($n = 10$ centers; 186
12 children), or 2) waiting list – control group ($n = 9$ centers; 175 children). Saliva samples for
13 cortisol levels were collected 3 times/day, pre- and post-implementation. The Minipally puppet
14 program consists of 2 workshops/month for 8 months for the development of social skills and
15 self-regulation in 2-5-year-olds, with reinforcement activities between workshops. Educators
16 received 2-days' training and 12 hours' supervision in Minipally.

17 **Results:** Linear mixed models for repeated measures revealed a significant interaction between
18 intervention status and time of day of cortisol sampling ($\beta = -0.18$, $p = 0.04$). The intervention
19 group showed patterns of decreasing diurnal cortisol secretion ($\beta = -0.32$, $p < 0.01$), whereas
20 the control group showed increasing slopes ($\beta = 0.20$, $p < 0.01$). Moreover, family income was a
21 moderator; children in lower-income families benefited most from the intervention.

22 **Conclusion:** Results suggest that a social skills training program, when integrated into a
23 preschool education curriculum, can foster an environment more conducive to typical childhood
24 patterns of cortisol secretion.

25

26 *Keywords:* Child care, daycare, social skills training, cortisol, diurnal cortisol, low income

1 *Abbreviations:* HPA, Hypothalamic-pituitary-adrenal.

2

3 **Trial registration number:** ISRCTN84339956 (retrospectively registered in March 2017, no
4 amendment to initial protocol).

5

1 Impact of a social skills program on children's stress: A cluster randomized trial

2 **1. Introduction**

3 In most Western industrialized countries, the percentage of preschool children receiving child
4 care services has increased dramatically since the mid-1980s (OECD, 2014). In this context, we
5 will use the term "child care" to designate a daycare center; i.e. a preschool or regular group-
6 based care of children prior to school entry by someone other than the parents, who are
7 generally at work. In the majority of countries belonging to the Organization for Economic Co-
8 operation and Development (OECD), it is estimated that at least 80% of children receive full-
9 time child care before they enter elementary school (OECD, 2014). Child care constitutes a
10 promising and supportive environment, especially for children from socioeconomically
11 disadvantaged neighborhoods, in at-risk families, or with early socioemotional maladjustment
12 (Côté et al., 2007; Herba et al., 2013).

13 For many children, child care represents the first social experience in a structured group
14 setting. Although child care is associated with a number of head start benefits for school
15 readiness (higher receptive vocabulary and reading skills), particularly for children from low
16 socioeconomic levels (Geoffroy et al., 2010), child care may also be a stressful experience,
17 especially for younger children or those attending lower-quality establishments (Ouellet-Morin et
18 al., 2010; Watamura et al., 2002). Two meta-analyses have reported that preschoolers (3–5
19 year-olds) tended to secrete higher diurnal levels of cortisol, the so-called "stress hormone",
20 when attending child care as compared to staying home (Geoffroy et al., 2006; Vermeer and
21 van Ijzendoorn, 2006). These flat or increasing patterns of cortisol secretion differ from the
22 gradually decreasing pattern biologically expected for children this age (Watamura et al., 2009).

23 *1.1 Child Care Center as a Stressful Environment*

24 The child care environment may be perceived by young children as unpredictable,
25 uncontrollable, and more threatening than their own home. Repeated and intense activation of

1 the hypothalamic-pituitary-adrenal (HPA) axis may lead to flattened or increasing diurnal cortisol
2 secretion (Gunnar et al., 2009). Although there is no evidence of a direct link between stress in
3 child care settings and poor health outcomes, chronic exposure to high levels of stress has
4 been shown to negatively impact child health (Koss and Gunnar, 2017). Furthermore, tolerable
5 and environmentally adaptive stress was shown to be associated to well-being (Gunnar and
6 Quevedo, 2007), warranting interventions that target potential sources of stress in child care
7 centers.

8 Two characteristics associated with the child care environment may induce stress and
9 activation of the HPA axis. First, developmentally, the age group in child care centers coincides
10 with peak levels of physically aggressive behaviors in response to conflict. This increases the
11 probability of victimization or of witnessing physical aggression (Côté et al., 2006). Indeed,
12 being either the perpetrator or the victim of physically aggressive behaviors may lead to high
13 levels of peer rejection, which in turn is also associated with rising cortisol patterns in children
14 attending child care (Gunnar and Donzella, 2002).

15 Second, preschoolers may not have reached sufficient levels of emotion regulation
16 (Gunnar and Donzella, 2002) to engage in prosocial behaviors. Emotion regulation has been
17 described as the ability to process emotions and emotional information, including inhibition of
18 emotional impulses, modulation of emotional behavior, and disengagement from distressing
19 elements (Grolnick & al., 2006). In children with poor emotion regulation skills, the risks of
20 disruptive behavior, poor social skills, and peer rejection are increased (Gunnar et al., 2003).

21 *1.2 Targeted Interventions for Children Attending Child Care Centers*

22 Results from relationship-focused interventions targeting children exposed to early life adversity,
23 such as foster care and neglect, suggest that improving the quality of the caregiver-child bond
24 may foster optimal HPA-axis activity (Bernard et al., 2015). To the best of our knowledge, only
25 one study tested the hypothesis that teacher-child relationships in child care could impact
26 cortisol levels (Hatfield and Williford, 2017). The objective of their 7-week program was to foster

1 sensitive teacher-child relationships and prosocial interactions among disruptive children (n =
2 113). Lower cortisol levels in the morning were observed for children exposed to the intervention
3 as compared to the control group. These results suggest that a psychosocial program within the
4 child care environment can modify HPA-axis activity.

5 *1.3 Targeted Interventions for Child Care Centers in Low-Income Neighborhoods*

6 Low-income neighborhoods present a confluence of distal and proximal challenges thought to
7 influence biological, cognitive, and behavioral development in children (Koss and Gunnar,
8 2017). Specifically, child care centers in low-income neighborhoods tend to be fewer and of
9 lower quality than in higher-income neighborhoods (Cloney et al., 2016). Also, children from
10 disadvantaged socioeconomic backgrounds are more likely to have cognitive and
11 socioemotional deficits (Cloney et al., 2016; Goldfeld et al., 2015) which may lead to higher
12 levels of aggression in the child care setting. However, children from lower-income families also
13 stand to benefit the most from attending good quality child care, especially in terms of social
14 development (Côté et al. 2007). Interventions supporting the development of prosocial skills and
15 emotion regulation in low-income neighborhood child care centers may lower diurnal cortisol
16 secretion, by reducing the number of conflicts to which the children are exposed. Social skills
17 training might also buffer HPA-axis activation by lowering levels of peer exclusion, increasing
18 prosocial interactions with peers, and encouraging children to ask for help from a trusted
19 caregiver (Hostinar and Gunnar, 2015).

20 *1.4 Study Aims and Hypothesis*

21 We conducted a cluster randomized controlled trial of a training program for early childhood
22 educators to implement in child care centers of low-income neighborhoods. The aim was to test
23 whether preschoolers attending a child care center with a social skills intervention component,
24 by educators with specific training, showed more normative patterns of diurnal cortisol secretion
25 than children not exposed to such a program. Additionally, we tested whether the impact of the

1 intervention varied as a function of family income, whereby children from lower-income families
2 were hypothesized to benefit the most from social skills training.

3

4 **2. Methods**

5 *2.1 Participants*

6 We conducted the study in low-income neighborhoods of Montreal, Quebec, Canada from
7 September 2013 to June 2014. Eligibility criteria were that at least 25% of the children attending
8 the child care come from low-income families, defined as those entitled to a special government
9 subsidy providing free child care access to families with an annual family income below
10 Can\$20,000. The income status of the neighborhood itself was defined according to both
11 provincial (Québec., 2013) and national criteria (Canada, 2011). Children were not clustered in
12 particular child care centers by familial income and the distribution of family income was not
13 uniform across child care centers, allowing us to investigate the moderating effect of family
14 income on intervention status and cortisol secretion. Of 38 child care centers manifesting an
15 interest, 19 were eligible. We determined that a sample of 19 centers would allow us to detect a
16 small-to-medium effect size, with 90% power at a two-sided significance level of 5% (Heo,
17 2008).

18 *2.2 Study Design*

19 Randomization was at the level of the child care center. Centers were randomized to receive the
20 Minipally social skills program in Year 1 (intervention group) or Year 2 (waiting list, control
21 group). Each child care center included 2-3 classes of 4-year-olds, with 8 children per class.
22 Altogether, 43 classes ($n = 361$ children) in 19 child care centers took part in the study (Figure
23 1). Cluster randomization ensured that children from the control (waiting list) group were not
24 exposed to the intervention. The intervention team provided Minipally training to the educators,
25 and the research team evaluated the impact of the intervention. Once data collection was
26 complete, all child care centers in the control group received the Minipally training. Consent to

1 participate in the study was obtained from parents and child care directors. The Sainte-Justine
2 Hospital Research Ethics Board approved all procedures in May 2013. A detailed description of
3 the study protocol has been published (Côté et al., 2017).

4 Insert Figure 1 here

5 *2.3 Intervention*

6 The intervention was the Minipally program for the development of social skills and self-
7 regulation in children aged 2 to 5 years. The Minipally curriculum is delivered via a puppet that
8 presents itself as a loyal and enthusiastic friend visiting the child care to model prosocial
9 behaviors and social inclusion. There are 16 play sessions (2/month for 8 months) where the
10 educators and puppet discuss/play with friends (other puppets) and the children. The program
11 includes generic components of social skills training for children: introduction to social contact
12 (make and accept contact from others, make requests); problem-solving (identifying the
13 problem, generating solutions); self-regulation (breathing to calm down, accepting frustration,
14 learning to share, tolerating frustration); and emotion regulation (identifying and expressing
15 emotions, listening to others).

16 Child care educators at centers randomized to the intervention received 2 days of
17 intensive training in the Minipally program plus 12 hours of classroom supervision (4 half-day
18 sessions over the course of the program). After each Minipally session with the children,
19 educators reinforced the principles with 2 weeks of activities based on the topic addressed (for
20 instance, if the Minipally puppet intervened in a particular way in a conflict between two children,
21 then over the next 2 weeks, the educator designed activities to recall the strategies presented
22 by the puppet).

23 **3. Measures**

24 *3.1 Saliva Collection*

25 Saliva samples were collected before and after the intervention by trained research assistants
26 blinded to the status of the child care center (intervention or control). Samples were collected at

1 3 time points: 1) 30 minutes after the child's arrival at child care (between 7:00 and 10:30 a.m.);
2 2) before lunch (between 10:45 a.m. and 12:00 a.m.); and 3) one hour after waking from the
3 afternoon nap (between 2:30 and 4 p.m.). Saliva was collected by placing a cotton sponge
4 under the child's tongue for one minute (SalivaBio Children's Swab, Salimetrics).

5 Samples were stored at -20°C until the laboratory assays. Cortisol concentration was
6 obtained using a high-sensitivity enzyme immunoassay where the lowest limit of detection was
7 0.007µg/dL. All samples were assayed in duplicate. The intra- and inter-assay coefficients of
8 variation were 4.46% and 8.28%, respectively.

9 *3.2 Parent Questionnaires*

10 Parents answered a short questionnaire upon arrival at the child care center, both before and
11 after the intervention. Questionnaires included a wide range of variables potentially affecting
12 cortisol secretion, such as children's habits and waking time, foods eaten for breakfast, current
13 medication, sleep quality the night before, current mood (sad, excited), and state of health (cold,
14 allergies).

15 Information on sociodemographic background (parent education and family income) was
16 collected before the intervention. Annual family income ranged from "less than Can\$10,000" to
17 "more than Can\$80,000." Because the income distribution was not uniform, and to ensure a
18 similar number of children in each group, we split the sample into three: lower-income families
19 entitled to government subsidy (income less than Can\$20,000), middle-income families
20 (Can\$20,000 to Can\$80,000), and higher-income families (over Can\$80,000).

21 *3.3 Children's Social Behaviors Assessed by Child Care Educators*

22 Educators completed the social behavioral questionnaire (Tremblay et al., 1992) for each child
23 in their group at pre- and post-intervention. Two dimensions of the validated and well-published
24 questionnaire (Pingault et al., 2011) were used: a) disruptive behaviors including five items on
25 opposition (e.g., has been defiant or has refused to comply with an adult request); four on
26 impulsivity/hyperactivity questions (e.g., has had difficulty waiting for his/her turn in games); six

1 on physical aggression questions (three reactive, e.g., has reacted in an aggressively when
2 teased, and three non-reactive, e.g., has gotten into fights) (Cronbach alpha = 0.86); and b)
3 prosocial behaviors (e.g., has helped other children; 7 items) (Cronbach alpha = 0.79).

4 *3.4 Child Care Educator Questionnaires*

5 Every educator completed a sociodemographic questionnaire as well as a questionnaire
6 assessing their training in early childhood education and care. Educators in the intervention
7 group also completed a logbook in which they indicated when and which Minipally activities had
8 been conducted in their group.

9 **4. Data Analysis**

10 *4.1 Preliminary analyses*

11 We compared the intervention and control groups at baseline for 35 variables that might directly
12 or indirectly affect the impact of the intervention, including age, family income and number of
13 hours of child care per week. Less than 6% of the variables differed between groups,
14 suggesting that randomization was successful. Nonetheless, we verified whether those
15 variables were associated with diurnal cortisol secretion. None were, so we did not control for
16 them in subsequent models.

17 Second, we identified outlier values for which the cortisol concentration was above 3
18 standard deviations of the sample mean. At each saliva sample collection time, there were 2-5
19 outliers (for a total of 23); these were winsorized so as not to exert undue influence on the
20 results. We then performed a square root transformation of the cortisol distribution at each time
21 point to account for skewness. We used bivariate analyses (Spearman correlation coefficients
22 and ANOVA) to search for potential covariates associated with cortisol levels, such as exact
23 hour of sampling. For each cortisol sample, we modeled multiple variable linear regressions
24 using the following independent variables: exact hour of sampling, time of awakening, hours
25 since last saliva sample and all potential confounders related to cortisol secretion. We derived
26 the residuals from each linear regression (i.e. one regression for every cortisol sample) so that

1 the main analyses would be free from potential confounders. Raw cortisol concentration and
2 residual concentration at each sample time according to intervention groups are presented in
3 Table S3 in Supplementary material.

4 *4.2 Main Analyses*

5 To test the impact of the intervention on diurnal cortisol secretion (i.e. cortisol change during the
6 day), we used linear mixed models for repeated measures. This method allows the modeling of
7 multiple data points nested within individuals while also modeling between-subject differences.
8 Accounting for non-independence of repeated measures, linear mixed modeling allowed for the
9 possibility that cortisol samples taken on the same day in any given child might be more
10 correlated than samples in different children. Each child was assigned an individual intercept
11 and slope of diurnal cortisol secretion. Patterns were adjusted for missingness. The final model
12 represents the mean effect of every intercept and slope (i.e. diurnal cortisol secretion) according
13 to intervention status (intervention vs. control). We did not include an additional level for child
14 care as the intracluster correlation coefficient for child care was less than 5%. Analyses with and
15 without this level yielded similar results, so we reported the most parsimonious model
16 (Tabachnick and Fidell, 2007).

17 Linear mixed modeling was conducted pre-and post-intervention. At each time point, the
18 fixed effects were intervention status (intervention vs. control), time of saliva collection (morning
19 [0], just before lunch [1], and one hour after waking from the afternoon nap [2]), and interaction
20 (intervention status x time of saliva collection). At post-intervention, we also controlled for pre-
21 intervention cortisol levels. We also examined whether the effects of the intervention varied
22 according to family income. More specifically, we tested a three-way interaction including
23 intervention status, time of saliva collection, and family income. All analyses were conducted
24 using IBM SPSS Statistics, version 24 (Armonk, NY, USA).

25

26 **5. Results**

1 sample ($F = 27.34$, $df = 114$, $p < 0.01$) where children from low and middle-income families had
2 their lowest level of diurnal cortisol concentration.

3 *5.4 Supplementary analyses.*

4 To explore the possibility that the impact of the intervention occurred via changes in child
5 behaviors, we tested whether disruptive and prosocial behaviors assessed by child care
6 educators in post-intervention mediated the effect of the intervention on cortisol secretion.
7 Specifically, we used two formulas for the calculation of overall diurnal cortisol secretion by
8 areas under the curve: 1) the “Area under the curve with respect to increase” (AUCI) and “Area
9 under the curve with respect to ground” (AUCG) (Pruessner et al., 2003). Using these
10 outcomes, intervention status was associated with AUCG ($\beta = -1.62$, $p = 0.01$), and not with
11 AUCI ($\beta = -1.32$, $p = 0.07$). The intervention was also associated with a significant decrease of
12 children’s disruptive behaviors in post-intervention ($\beta = -0.61$, $p = 0.02$), but disruptive behaviors
13 were not associated with children’s AUCG ($\beta = -0.09$, $p = 0.52$) or AUCI ($\beta = 0.03$, $p = 0.85$).
14 The intervention was not associated with an increase of children’s prosocial behaviors in post-
15 intervention ($\beta = 0.38$, $p = 0.09$), and prosocial behaviors were not associated with children’s
16 AUCG ($\beta = 0.01$, $p = 0.99$) or AUCI ($\beta = 0.01$, $p = 0.63$). Thus, neither disruptive nor prosocial
17 behaviors in post-intervention were predictors of overall cortisol secretion. Finally, we
18 investigated if family income was also a moderator of the association between intervention
19 status and children’s social behaviors in the mediation models between intervention status and
20 children’s overall diurnal cortisol secretion. We did not find a significant moderation by familial
21 income for disruptive behaviors ($p = 0.30$) neither for prosocial behaviors ($p = 0.65$).

22

23

Insert Table 2 here

24

Insert Figure 3 here

25

1 **6. Discussion**

2 Young children in child care have been reported to exhibit flat levels of diurnal cortisol secretion,
3 as opposed to the decreasing pattern expected for children that age (Vermeer and van
4 Ijzendoorn, 2006; Watamura et al., 2002). We conducted a cluster randomized controlled trial to
5 test whether a social skills training program, led by early childhood educators in child care
6 centers, would restore expected patterns. Whereas both the intervention and control groups in
7 participating child care centers exhibited flat diurnal cortisol secretion at baseline, we found that
8 children exposed to the intervention exhibited more typical, decreasing levels of cortisol
9 secretion throughout the day, as compared to controls. Further, children from lower-income
10 families seemed to benefit more from the intervention than those from middle- or higher-income
11 backgrounds. Interestingly, in the intervention group, children from higher-income families had
12 diurnal patterns characterized by early suppression of morning cortisol, whereas children from
13 middle- and low-income families had greater cortisol suppression in the afternoon, as compared
14 to controls.

15 A possible explanation for the sharper decline in the diurnal cortisol slope in children
16 from lower-income families is that they may be more sensitive than others to the intervention
17 and to the ensuing changes in the child care environment. Children from disadvantaged
18 backgrounds are more likely to be disruptive (Shaw and Shelleby, 2014) and to be victimized in
19 peer-play interactions (Barker et al., 2008). The social skills intervention might have enhanced
20 the dynamic of the group as a whole, thereby helping the children most at risk. Fewer social
21 challenges and confrontations in the intervention group might explain the observed cortisol
22 decreases. For the control group, the increasing cortisol levels observed for low- and middle-
23 income children similarly point to the idea that these children were in need of an intervention
24 fostering a less stressful environment.

25 A complementary hypothesis is that children from higher-income families might be more
26 sensitive to age-appropriate social challenges in the child care setting, as compared to children

1 from middle- or lower-income families, assuming that children from higher-income families faced
2 fewer challenges in their home environment (Ellis and Boyce, 2008). For children of higher-
3 income families in both the intervention and control groups, the child care environment might
4 still be perceived as unpredictable and uncontrollable, two features associated with greater
5 cortisol response (Gunnar et al., 2009). The diurnal patterns observed in this study support the
6 idea that HPA-axis activation in child care is an adaptive reaction to the perception of challenge
7 and unpredictability in the environment. More generally, these findings support the concept that
8 children exposed to different levels of adversity will perceive threats differently and react to them
9 differently in daily life (Ellis and Del Giudice, 2014).

10 Finally, we note that our results are in line with those reported by Berry and colleagues
11 (Berry et al., 2016; Berry et al., 2014), where children in child care from lower socioeconomic
12 status exhibited lower levels of diurnal cortisol than their wealthier peers. However, despite
13 similarities between our results and those obtained in the Family Life Project (Berry et al., 2016;
14 Berry et al., 2014), the mechanisms underlying interactions between family income and diurnal
15 cortisol secretion remain largely unclear and there is some not-yet-understood complexity in the
16 present results.

17 To the best of our knowledge, there is no longitudinal study examining the impact of high
18 stress levels on later physical and mental health in children who regularly attend child care. We
19 are therefore limited in the discussion of potential long-term effects of our intervention on well-
20 being and development. Future studies should investigate the consequences of flattened and
21 increasing diurnal cortisol secretion in children attending child care, as it is a developmental and
22 context-specific phenomenon that has been replicated several times in different welfare systems
23 (Geoffroy et al., 2006; Vermeer and van Ijzendoorn, 2006).

24 It is important to emphasize three features of the social skills program evaluated in this
25 study. First, the program is inclusive—all children are involved in the social skills training—not
26 only those exhibiting socioemotional or behavioral problems. This lowers the total cost of the

1 intervention and notably, reduces the risk of stigma in children. Second, the program can easily
2 be integrated into daily routine and educational activities, thus increasing adherence levels of
3 early childhood educators to the program. Third, the program can be disseminated with
4 relatively low implementation costs and educator training. Dividing the total expenses for
5 program implementation (educator training, supervision, and monetary compensation to child
6 care directors) by the number of children who received the intervention ($n = 185$), the total
7 cost/child works out to Can\$95.

8 It is noteworthy that we did not find an association between social behaviors (either
9 disruptive or prosocial) and diurnal cortisol secretion in supplementary analyses. Further
10 research will be needed to address the potential effect of social skills training on peer-peer
11 relationships, educator-child relationship, the degree of classroom chaos, and the social
12 behaviors of the group to identify the mechanisms involved in the interaction between the
13 intervention and HPA-axis activation.

14 *6.1 Strengths and Limitations*

15 The main strengths of this study are its cluster-randomized experimental design and the
16 measurement of diurnal cortisol levels three times per day to determine secretion patterns.
17 Further, cortisol samples were collected by our research assistants, ensuring that collection
18 procedures such as sampling time were uniform throughout. The study has high ecological
19 validity; it was conducted in community-based child care centers, with early childhood educators
20 whose qualifications were the two-year college program in early childhood and child care
21 education required by the Quebec government (as compared to fully licensed psychologists,
22 teachers, or social workers). Finally, our sample was sufficiently heterogeneous to justify testing
23 family income as a moderator between intervention status and diurnal cortisol secretion.

24 The study also had some limitations. First, 6% of the children left child care over the
25 course of the study. While for the most part, the family sociodemographic characteristics of
26 newcomers were no different from those who left, we could not statistically control for pre-

1 intervention cortisol secretion in children who joined the study later. Second, we do not know the
2 exact number of workshops animated by child care educators. We only know that 90% of the
3 educators performed 12 or more workshops out of 16 during the year of implementation. Future
4 studies should include comprehensive implementation evaluation. Third, as we did not collect
5 longer-term post-intervention cortisol samples, we did not test whether the reported differences
6 persisted over time. Similarly, we did not test whether the observed decline in cortisol secretion
7 was associated with a lower risk of behavioral difficulties later on. Such investigation would have
8 required additional time points to test this hypothesis according to a clear temporal sequence
9 where the HPA axis would be a mediator between the psychosocial intervention targeting
10 quality child care and disruptive behavior. Replication of this study with long-term follow-up,
11 larger sample size, and different models of child care (center-based vs. home-based) is needed
12 to advance knowledge on the mechanisms linking child care quality to diurnal patterns of
13 cortisol secretion and long-term social and emotional development.

14

15 **7. Conclusion**

16 Our findings suggest that a social skills training program, when integrated into a preschool
17 education curriculum, can foster an environment more conducive to typical childhood patterns of
18 cortisol secretion. Given that child care services cover a critical developmental period during
19 which young children learn to interact in a group setting, this program may be a promising way
20 to ensure and promote health and well-being from an early age.

21

22 ***Acknowledgements***

23 The authors wish to thank Danielle Buch, medical writer, for critical revision and substantive
24 editing of the manuscript. We are grateful to the participants who have given their time to take
25 part in this study.

1

2 **Figure Legends**

3 **Figure 1.** Trial Flow Diagram

4

5 **Figure 2.** Post-intervention diurnal cortisol secretion, according to intervention status

6 Abbreviations: AM, morning; PM, afternoon.

7 Note: Post-intervention refers to 8 months after the start of the intervention.

8

9 **Figure 3.** Post-intervention diurnal cortisol secretion, according to family income and

10 intervention status

11 Abbreviations: AM, morning; PM, afternoon.

12 Note 1. Post-intervention refers to 8 months after the start of the intervention.

13 Note 2. Cortisol concentration residuals were derived from the three-way interaction between
14 family income, intervention status, and time of day.

15

16 **Supplementary Material**

17 **Figure S4.** Post-intervention diurnal cortisol secretion, according to intervention status and

18 family income

19 Abbreviations: AM, morning; PM, afternoon.

20 Note 1. Post-intervention refers to 8 months after the start of the intervention.

21 Note 2. Cortisol concentration residuals were derived from the three-way interaction between
22 family income, intervention status, and time of day.

23

1 **References**

- 2 Barker, E.D., Boivin, M., Brendgen, M., Fontaine, N., Arseneault, L., Vitaro, F., Bissonnette, C.,
3 Tremblay, R.E., 2008. Predictive validity and early predictors of peer-victimization trajectories in
4 preschool. *Archives of General Psychiatry* 65, 1185-1192.
- 5 Bernard, K., Hostinar, C.E., Dozier, M., 2015. Intervention effects on diurnal cortisol rhythms of
6 Child Protective Services-referred infants in early childhood: preschool follow-up results of a
7 randomized clinical trial. *JAMA Pediatr* 169, 112-119.
- 8 Berry, D., Blair, C., Granger, D.A., Investigators, F.L.P.K., 2016. Child care and cortisol across
9 infancy and toddlerhood: poverty, peers, and developmental timing. *Family relations* 65, 51-72.
- 10 Berry, D., Blair, C., Ursache, A., Willoughby, M., Garrett-Peters, P., Vernon-Feagans, L.,
11 Bratsch-Hines, M., Mills-Koonce, W.R., Granger, D.A., Family Life Project Key, I., 2014. Child
12 care and cortisol across early childhood: context matters. *Dev Psychol* 50, 514-525.
- 13 Canada, S., 2011. *Census and 2011 National Household Survey*. Statistic Canada, Canada.
- 14 Cloney, D., Cleveland, G., Hattie, J., Tayler, C., 2016. Variations in the availability and quality of
15 early childhood education and care by socioeconomic status of neighborhoods. *Early Education
16 and Development* 27, 384-401.
- 17 Côté, Boivin, M., Nagin, D.S., Japel, C., Xu, Q., Zoccolillo, M., Junger, M., Tremblay, R.E.,
18 2007. The role of maternal education and nonmaternal care services in the prevention of
19 children's physical aggression problems. *Archives of general psychiatry* 64, 1305-1312.
- 20 Côté, Larose, M.-P., Geoffroy, M.C., Laurin, J., Vitaro, F., Tremblay, R.E., Ouellet-Morin, I.,
21 2017. Testing the impact of a social skill training versus waiting list control group for the
22 reduction of disruptive behaviors and stress among preschool children in child care: the study
23 protocol for a cluster randomized trial. *BMC psychology* 5, 29.
- 24 Côté, S., Vaillancourt, T., LeBlanc, J.C., Nagin, D.S., Tremblay, R.E., 2006. The development of
25 physical aggression from toddlerhood to pre-adolescence: A nation wide longitudinal study of
26 Canadian children. *Journal of abnormal child psychology* 34, 68-82.

- 1 Ellis, B.J., Boyce, W.T., 2008. Biological sensitivity to context. *Current Directions in*
2 *Psychological Science* 17, 183-187.
- 3 Ellis, B.J., Del Giudice, M., 2014. Beyond allostatic load: Rethinking the role of stress in
4 regulating human development. *Development and Psychopathology* 26, 1-20.
- 5 Geoffroy, M.C., Cote, S.M., Giguere, C.E., Dionne, G., Zelazo, P.D., Tremblay, R.E., Boivin, M.,
6 Seguin, J.R., 2010. Closing the gap in academic readiness and achievement: the role of early
7 childcare. *J Child Psychol Psychiatry* 51, 1359-1367.
- 8 Geoffroy, M.C., Cote, S.M., Parent, S., Seguin, J.R., 2006. Daycare attendance, stress, and
9 mental health. *Can J Psychiatry* 51, 607-615.
- 10 Goldfeld, S., Woolcock, G., Katz, I., Tanton, R., Brinkman, S., O'Connor, E., Mathews, T., Giles-
11 Corti, B., 2015. Neighbourhood effects influencing early childhood development: conceptual
12 model and trial measurement methodologies from the Kids in Communities Study. *Social*
13 *Indicators Research* 120, 197-212.
- 14 Gunnar, Donzella, B., 2002. Social regulation of the cortisol levels in early human development.
15 *Psychoneuroendocrinology* 27, 199-220.
- 16 Gunnar, M.R., Seban, A.M., Tout, K., Donzella, B., van Dulmen, M.M., 2003. Peer rejection,
17 temperament, and cortisol activity in preschoolers. *Developmental Psychobiology: The Journal*
18 *of the International Society for Developmental Psychobiology* 43, 346-368.
- 19 Gunnar, M.R., Talge, N.M., Herrera, A., 2009. Stressor paradigms in developmental studies:
20 What does and does not work to produce mean increases in salivary cortisol.
21 *Psychoneuroendocrinology* 34, 953-967.
- 22 Hatfield, B.E., Williford, A.P., 2017. Cortisol Patterns for Young Children Displaying Disruptive
23 Behavior: Links to a Teacher-Child, Relationship-Focused Intervention. *Prevention Science* 18,
24 40-49.
- 25 Heo, M., & Leon, A. C. Statistical power and sample size requirements for three level
26 hierarchical cluster randomized trials. *Biometrics*, 64(4), 1256-1262., 2008. Statistical power and

- 1 sample size requirements for three level hierarchical cluster randomized trials. . *Biometrics* 64,
2 1256-1262.
- 3 Herba, C.M., Tremblay, R.E., Boivin, M., Liu, X., Mongeau, C., Seguin, J.R., Cote, S.M., 2013.
4 Maternal depressive symptoms and children's emotional problems: can early child care help
5 children of depressed mothers? *JAMA Psychiatry* 70, 830-838.
- 6 Hostinar, C.E., Gunnar, M.R., 2015. Social support can buffer against stress and shape brain
7 activity. *AJOB neuroscience* 6, 34-42.
- 8 Koss, K.J., Gunnar, M.R., 2017. Annual Research Review: Early adversity, the hypothalamic–
9 pituitary–adrenocortical axis, and child psychopathology. *Journal of Child Psychology and*
10 *Psychiatry*.
- 11 OECD, 2014. PF3.2: Enrolment in childcare and pre-schools. , Paris: France.
- 12 Ouellet-Morin, I., Tremblay, R.E., Boivin, M., Meaney, M., Kramer, M., Cote, S.M., 2010. Diurnal
13 cortisol secretion at home and in child care: a prospective study of 2-year-old toddlers. *J Child*
14 *Psychol Psychiatry* 51, 295-303.
- 15 Pingault, J.-B., Tremblay, R.E., Vitaro, F., Carbonneau, R., Genolini, C., Falissard, B., Côté,
16 S.M., 2011. Childhood trajectories of inattention and hyperactivity and prediction of educational
17 attainment in early adulthood: a 16-year longitudinal population-based study. *American Journal*
18 *of Psychiatry* 168, 1164-1170.
- 19 Pruessner, J.C., Kirschbaum, C., Meinlschmid, G., Hellhammer, D.H., 2003. Two formulas for
20 computation of the area under the curve represent measures of total hormone concentration
21 versus time-dependent change. *Psychoneuroendocrinology* 28, 916-931.
- 22 Québec., R.d.r.d., 2013. Child Assistance Program. Ministère du Revenu du Québec.
- 23 Shaw, D.S., Shelleby, E.C., 2014. Early-starting conduct problems: Intersection of conduct
24 problems and poverty. *Annual Review of Clinical Psychology* 10, 503-528.
- 25 Tabachnick, B.G., Fidell, L.S., 2007. Using multivariate statistics. Allyn & Bacon/Pearson
26 Education.

- 1 Tremblay, R.E., Vitaro, F., Gagnon, C., Piché, C., Royer, N., 1992. A prosocial Scale for the
2 Preschool Behaviour Questionnaire: Concurrent and Predictive Correlates. *International Journal*
3 *of Behavioral Development* 15, 227-245.
- 4 Vermeer, H.J., van Ijzendoorn, M.H., 2006. Children's elevated cortisol levels at daycare: A
5 review and meta-analysis. *Early Childhood Research Quarterly* 21, 390-401.
- 6 Watamura, S.E., Kryzer, E.M., Robertson, S.S., 2009. Cortisol patterns at home and child care:
7 Afternoon differences and evening recovery in children attending very high quality full-day
8 center-based child care. *Journal of Applied Developmental Psychology* 30, 475-485.
- 9 Watamura, S.E., Sebanc, A.M., Gunnar, M.R., 2002. Rising cortisol at childcare: relations with
10 nap, rest, and temperament. *Dev Psychobiol* 40, 33-42.

11

12

Figure 1. Trial Flow Diagram





Table 1. Children's descriptive statistics at baseline

	Control group (n = 175)	Intervention group (n = 186)	<i>P</i> value
Age, months ^a	52.8 (5.0)	54.5 (4.5)	< 0.01
Sex (<i>boy</i>) ^b	86 (49.1%)	98 (52.7%)	0.57
	164	165	
Have siblings ^b	128 (78.5%)	143 (86.7%)	0.12
	163	165	
Child care hours/week ^b			
< 30 hours	24 (14.6%)	35 (21.2%)	0.26
30-40 hours	104 (63.4%)	95 (57.6%)	
> 40 hours	36 (22.0%)	35(21.2%)	
	164	165	
Family income ^b			
< Can\$19 999	34 (21.7%)	23 (14.7%)	0.26
Can\$20 000-80 000	69 (43.9%)	71 (45.5%)	
> Can\$80 000	54 (34.4%)	62 (39.7%)	
	157	156	
Highest maternal education ^b			
High school diploma	20 (12.7%)	21 (13.1%)	0.53
Vocational training	49 (31.0%)	48 (30.0%)	
Bachelor's degree	67 (42.4%)	60 (37.5%)	
Master's or PhD	22 (13.9%)	31 (19.4%)	
	158	160	

^a Mean (SD).

^b n (%).

Figure 2. Post-intervention diurnal cortisol secretion, according to intervention status



Abbreviations: AM, morning; PM, afternoon.


Note: Post-intervention refers to 8 months after the start of the intervention.

Table 2. Post-intervention diurnal cortisol secretion, according to family income and intervention status

	β (95% CI)	<i>P</i> value
Intercept	-0.63 (-1.08, -0.17)	0.01
Pre-intervention cortisol levels	0.18 (0.11, 0.26)	< 0.01
Intervention	0.53 (-0.16, 1.23)	0.13
Time	0.38 (-0.01, 0.76)	0.05
Time x intervention	-0.79 (-1.36, -0.22)	0.01
Income	0.26 (0.06, 0.45)	0.01
Income x intervention	-0.23 (-0.53, 0.07)	0.13
Income x time	-0.14 (-0.30, 0.02)	0.09
Income x intervention x time	0.27 (0.02, 0.51)	0.03

Abbreviation: CI, confidence interval.

Figure 3. Post-intervention diurnal cortisol secretion, according to family income and intervention status



Abbreviations: AM, morning; PM, afternoon.

Note 1. Post-intervention refers to 8 months after the start of the intervention.

Note 2. Cortisol concentration residuals were derived from the three-way interaction between family income, intervention status, and time of day.